

### Clarification Regarding “Outdoor Transmission of SARS-CoV-2 and Other Respiratory Viruses: A Systematic Review”

TO THE EDITOR—On April 24, 2021, the Centers for Disease Control and Prevention issued new guidance on outdoor activities [1]. In subsequent testimony before the US Senate, a number reported in our article, “Outdoor Transmission of SARS-CoV-2 and Other Respiratory Viruses: A Systematic Review,” was cited that the proportion of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) transmission occurring in outside settings is less than 10% [2].

We are writing to clarify how we arrived at the less than 10% summary number. Our abstract and results sections state that “five identified studies found a low proportion of reported global SARS-CoV-2 infections occurred outdoors (<10%).” Because of the small number of heterogeneous studies we reviewed, as well as their methodological limitations, we could not provide a meta-analytically pooled estimate of the exact proportion of SARS-CoV-2 transmissions that have occurred outdoors or the associated risk.

Ten percent was chosen as a conservative estimate based on the upper confidence limit of the proportion of cases attributable to outdoor settings reported in one of the studies we reviewed. The Lan et al [3] study, a work-related outbreak in Asia, reported that 5 (5%) of 103 cases were linked to construction sites; 10% is the upper limit of transmission potentially linked to an outdoor setting from this study. However, this number does not reflect the potential for community-acquired infection, nor does it accurately classify the total person-time at risk for transmission indoors versus outdoors in this occupational group. The other studies included in our review were

Qian et al [4] (<1%, or 2 of 7324 cases occurred outdoors), Nishiura et al [5] (odds of indoor transmission 18 times higher than outdoors, no raw data available) [5], Leclerc et al [6] (<1% transmissions or 95 of 10 926 total cases reported), and Szablewski et al [7], which described an outbreak of SARS-CoV-2 during a summer camp. Given the paucity and limitations of identified data and timing of study (ie, still early in pandemic), we reported 10%—the highest upper confidence interval in any of the studies we reviewed—to signify a small risk when compared with indoor settings, but not so negligible that it should be ignored pending further studies that could help clarify the risk more accurately.

As we discussed, our review has several limitations. The reports we found may have underestimated actual infections because cases reported may have missed asymptomatic infections or infections not reported to state agencies. The interpretation of the findings of the articles we included also may have led to overestimation in the number of outdoor transmissions, because the definition of outdoors was heterogeneous, and that, for example, transmissions in construction sites and in camp settings may be related to indoor time and not outdoor time. Therefore, deducing risks attributable to any policy decision and impact on SARS-CoV-2 transmission based on these data is subject to the potential limitations of these studies. Furthermore, as in any systematic review based on peer-reviewed publications, our review is subject to publication bias—results of all outbreak investigations may not be systematically published in peer-reviewed journals.

Given the growing body of evidence pertaining to the risk of outdoor transmission and policy implication of our findings, we are in the process of updating our systematic review. Additional relevant studies found to date range in

study design and quality but augment our findings that the risk of respiratory virus transmission outdoors is much lower than indoors [8–10].

Among these is a study that examined low secondary transmission rates of SARS-CoV-2 infection among contacts of construction laborers in India: 496 close contacts of 18 SARS-CoV-2-infected construction laborers were assessed for infection and exposure time, both outdoors and indoors, at work as well as at home. The study reports a low secondary attack rate (1.4%) among contacts of construction laborers at locations with free air flow compared with a 30-fold higher rate of transmission among households. Their primary data may help further delineate the incidence rate ratio of outdoor versus indoor infections—a number that may be more useful than proportions in explaining the relative risk of infections outdoors by accounting for person-time of exposure [8].

These updates support our initial conclusion that the risk of SARS-CoV-2 transmission is much lower outdoors than indoors. The proportion of infections happening outdoors is likely much lower than 10%; and some of the studies we have reviewed thus far suggest that it is likely less than 1%.

#### Notes

**Potential conflicts of interest.** All authors: No reported conflicts of interest. All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest.

Nooshin Razani,<sup>1,2</sup> Mohsen Malekinejad,<sup>1</sup> and George W. Rutherford<sup>1,2</sup>

<sup>1</sup>Department of Epidemiology and Biostatistics, University of California, San Francisco, San Francisco, California, USA;

<sup>2</sup>Department of Pediatrics, University of California, San Francisco, San Francisco, California, USA

#### References

- Centers for Disease Control. Guidance for Operating Youth and

- Summer Camps During COVID-19. Available at: <https://www.cdc.gov/coronavirus/2019-ncov/community/schools-childcare/summer-camps.html>. Accessed 26 May 2021.
2. Senator Collins to CDC Director: “Conflicting, Confusing Guidance from Your Agency has Undermined Public Confidence”. Susan Collins United States Senator for Maine Press Release, Posted on Senator Collins’s. Available at: <https://www.collins.senate.gov/newsroom/senator-collins-cdc-director-“conflicting-confusing-guidance-your-agency-has-undermined”>. Accessed 26 May 2021.
  3. Lan FY, Wei CF, Hsu YT, Christiani DC, Kales SN. Work-related COVID-19 transmission in six Asian countries/areas: a follow-up study. *PLoS One* **2020**; 15:e0233588.
  4. Qian H, Miao T, Liu L, Zheng X, Luo D, Li Y. Indoor transmission of SARS-CoV-2. *Indoor Air* **2021**; 31:639–45.
  5. Nishiura H, Oshitani H, Kobayashi T, et al. Closed environments facilitate secondary transmission of coronavirus disease 2019 (COVID-19). *medRxiv* **2020**. doi:10.1101/2020.02.28.20029272.
  6. Leclerc QJ, Fuller NM, Knight LE, Funk S, Knight GM; CMMID COVID-19 Working Group. What settings have been linked to SARS-CoV-2 transmission clusters? *Wellcome Open Res* **2020**; 5:83.
  7. Szablewski CM, Chang KT, Brown MM, et al. SARS-CoV-2 transmission and infection among attendees of an overnight camp - Georgia, June 2020. *MMWR Morb Mortal Wkly Rep* **2020**; 69:1023–5.
  8. Sundar V, Bhaskar E. Low secondary transmission rates of SARS-CoV-2 infection among contacts of construction laborers at open air environment. *Germs* **2021**; 11:128–31.
  9. Jones B, Phillips G, Kemp SP, et al. SARS-CoV-2 transmission during team-sport: do players develop COVID-19 after participating in rugby league matches with SARS-CoV-2 positive players? *Br J Sports Med* **2021**. doi: 10.1136/bjsports-2020-103714.
  10. Fouda B, Tran H, Makram O, et al. Identifying SARS-CoV2 transmission cluster category: an analysis of country government database. *J Infect Public Health* **2021**; 14:461–467.

Received 26 May 2021; editorial decision 28 May 2021; accepted 2 June 2021; published online June 4, 2021.

Correspondence: Nooshin Razani, MD, 550 16th St., 2nd Floor, San Francisco, CA 94158, USA ([nooshin.razani@ucsf.edu](mailto:nooshin.razani@ucsf.edu)).

**The Journal of Infectious Diseases®** 2021;XX:1–2  
 © The Author(s) 2021. Published by Oxford University Press for the Infectious Diseases Society of America. All rights reserved. For permissions, e-mail: [journals.permissions@oup.com](mailto:journals.permissions@oup.com). DOI: 10.1093/infdis/jiab298